

Higher K-energy functionals and higher Futaki invariants

Ben Weinkove

Department of Mathematics
Columbia University, New York, NY 10027

1. Introduction

In 1986, Mabuchi [11] introduced the K-energy functional, which integrates the Futaki invariant [8] and whose critical points are metrics of constant scalar curvature. It was used by Bando and Mabuchi [2] to prove a uniqueness theorem for Kähler-Einstein metrics. The K-energy is strongly related to notions of stability in Geometric Invariant Theory and has been used by Tian [13, 14, 16] and Phong and Sturm [12], to give some results on the conjecture of Yau [17, 18] on the relationship between stability and the existence of Kähler-Einstein metrics. Tian [15, 16] has cast both the K-energy and the Futaki invariant in a more general setting using Bott-Chern forms and Donaldson functionals.

Higher K-energy functionals were defined by Bando and Mabuchi [3] and generalize the K-energy map to higher Chern classes of the manifold. They integrate higher Futaki invariants (see [3], [1], [9], [10]). This note presents these functionals with an emphasis on Bott-Chern forms. Two new formulas for the higher K-energy are given and the second K-energy is shown to be related to Donaldson's Lagrangian applied to metrics on the tangent bundle.

Let M be a compact complex manifold of complex dimension n with Kähler metric $g_{i\bar{j}}$. Let $\omega = \sqrt{-1}g_{i\bar{j}}dz^i \wedge d\bar{z}^{\bar{j}}$ be the corresponding Kähler form. Define

$$P(M, \omega) = \{\phi \in C^\infty(M) \mid g_{i\bar{j}} + \partial_i \partial_{\bar{j}} \phi > 0\}.$$

For ϕ in $P(M, \omega)$, let $\{\phi_t\}_{0 \leq t \leq 1}$ be a smooth path in $P(M, \omega)$ with $\phi_0 = 0$, $\phi_1 = \phi$. Write ω_t for $\omega + \sqrt{-1}\partial\bar{\partial}\phi_t$. Now define the k th K-energy functional by

$$M_{k,\omega}(\phi) = -(n - k + 1) \int_0^1 \int_M \dot{\phi}_t (c_k(\phi_t) - \mu_k \omega_t^k) \wedge \omega_t^{n-k} dt,$$

where $c_k(\phi_t)$ (also written $c_k(\omega_t)$) is the representative of the k th Chern class of M given by the metric ω_t , and μ_k is the number given by

$$\mu_k = \frac{1}{V} \int_M c_k(\omega) \wedge \omega^{n-k}, \quad \text{where } V = \int_M \omega^n.$$

The case $k = 1$ corresponds to the original K-energy as defined in [11].

Theorems 1 and 2 below were proved by Bando and Mabuchi [3] (also see [9] for a proof of Theorem 1). We present an alternative derivation using Bott-Chern forms, leading to new formulas for the higher K-energy (Lemma 5.1 and Corollary 5.2) as well as Theorem 3 below, which gives the second K-energy in terms of Donaldson's Lagrangian.

Theorem 1. $M_{k,\omega}(\phi)$ is independent of the choice of path $\{\phi_t\}$.

To define the higher Futaki invariants, let $c_k(\omega)$ and $Hc_k(\omega)$ be respectively the k th Chern form and its harmonic part. There is a real smooth $(k-1, k-1)$ form f_k such that

$$c_k(\omega) - Hc_k(\omega) = \sqrt{-1} \partial \bar{\partial} f_k.$$

Define the k th Futaki invariant, which acts on a holomorphic vector field X , by

$$\mathcal{F}_{k,\omega}(X) = \int_M \mathcal{L}_X f_k \wedge \omega^{n-k+1},$$

where $\mathcal{L}_X$ is the Lie derivative. The case $k = 1$ corresponds to the original Futaki invariant if $c_1(M) = \mu_1 \omega$, so that $Hc_1(\omega) = \mu_1 \omega$. It is shown in [1] (see also [9]) that $\mathcal{F}_{k,\omega}$ is independent of choice of metric in the Kähler class.

Now given a holomorphic vector field X , let Φ_t be the integral curve of $X_R = X + \bar{X}$. Then $\Phi_t : M \rightarrow M$ is holomorphic, and there exists a smooth path $\{\phi_t\}$ in $P(M, \omega)$ satisfying

$$\Phi_t^* \omega = \omega + \sqrt{-1} \partial \bar{\partial} \phi_t, \quad \text{and} \quad \int_M \phi_t \omega^n = 0.$$

Then the higher K-energy integrates the Futaki invariant in the following sense.

Theorem 2. Suppose that $c_k(M) = \mu_k [\omega^k]$. Then with $\{\phi_t\}$ as above,

$$\frac{d}{dt} M_{k,\omega}(\phi_t) = 2 \operatorname{Re}(\mathcal{F}_{k,\omega}(X)).$$

Donaldson [6, 7] defined a Lagrangian which was used in his proof that Mumford-Takemoto stable vector bundles over projective algebraic varieties admit Hermitian-Einstein metrics. It is defined in terms of the Bott-Chern forms corresponding to $\text{Tr}(A)$ and $\text{Tr}(AB)$. It is unsurprising then that this Lagrangian is related to the first and second K-energy functionals, which correspond to the first and second Chern classes.

Let E be a holomorphic vector bundle of rank r over M , and let H and H_0 be Hermitian metrics on E . Let $\{H_t\}_{0 \leq t \leq 1}$ be a smooth path of metrics between H_0 and $H_1 = H$. Let F_t be the curvature of H_t . Then Donaldson's Lagrangian is given by

$$\begin{aligned} L(H, H_0) &= \int_M \int_0^1 n \sqrt{-1} \text{Tr}(\dot{H}_t H_t^{-1} F_t) \wedge \omega^{n-1} dt - \lambda \int_M \int_0^1 \text{Tr}(\dot{H}_t H_t^{-1}) \omega^n dt, \end{aligned}$$

where

$$\lambda = \frac{2\pi n}{r} \frac{1}{V} \int_M c_1(E) \wedge \omega^{n-1}.$$

Now consider the case when $E = T'M$, the holomorphic tangent bundle. Let H_0 be the Kähler metric g , and, for ϕ in $P(M, \omega)$, let H be g_ϕ . Write ω_ϕ for $\omega + \sqrt{-1} \partial \bar{\partial} \phi$ and $L(\phi)$ for $L(g_\phi, g)$. We have the following formula for the second K-energy functional.

Theorem 3. For ϕ in $P(M, \omega)$,

$$\begin{aligned} M_{2,\omega}(\phi) &= -\frac{1}{(2\pi)^2 n} L(\phi) \\ &\quad - \frac{1}{4\pi n} \int_M \log \left(\frac{\omega_\phi^n}{\omega^n} \right) (2\mu_1 \omega - nc_1(\phi) - nc_1(0)) \wedge \omega^{n-1} \\ &\quad - \int_M \phi \left(c_2(\phi) \wedge \sum_{i=0}^{n-2} \omega^i \wedge \omega_\phi^{n-2-i} - \left(\frac{n-1}{n+2} \right) \mu_2 \sum_{i=0}^n \omega^i \wedge \omega_\phi^{n-i} \right). \end{aligned}$$

I am very grateful to my advisor, D.H. Phong, for suggesting to look at these functionals, and for his subsequent guidance. I would also like to thank all the members of the Geometry and Analysis community at Columbia University for many enlightening discussions and talks. This paper will form a part of my forthcoming PhD thesis at Columbia University.

2. Chern classes and Bott-Chern forms

Before proving the above theorems, we will briefly discuss Chern classes and their Bott-Chern forms. For more details on Bott-Chern forms, see [4], [6] and [16]. Let E be a holomorphic vector bundle of rank r over the compact Kähler manifold M . If $H = H_{\alpha\bar{\beta}}$ is a Hermitian metric on E , then its curvature is an endomorphism-valued $(1,1)$ form given by

$$(F_H)_\alpha^\beta = (F_H)_\alpha^{\bar{i}\bar{j}} dz^i \wedge d\bar{z}^{\bar{j}} = \bar{\partial}((\partial H_{\alpha\bar{\gamma}}) H^{\bar{\gamma}\beta}).$$

Let Φ be an invariant symmetric k -linear function on $\mathfrak{gl}(r, \mathbf{C})$. Then we have a Chern-Weil form

$$\Phi(H) = \Phi(F_H) = \Phi(F_H, \dots, F_H) \in \Omega_M^{k,k},$$

which represents a characteristic class.

Let H and H_0 be two Hermitian metrics on E , and let $\{H_t\}$ be a smooth path of metrics between them. Then there exists a Bott-Chern form $\text{BC}_\Phi(H, H_0)$ in $\Omega_M^{k-1, k-1}/(\text{Im } \partial + \text{Im } \bar{\partial})$ given by

$$\text{BC}_\Phi(H, H_0) = -k\sqrt{-1} \int_0^1 \Phi(\dot{H}_t H_t^{-1}, F_{H_t}, \dots, F_{H_t}) dt.$$

It is shown in [4] that this definition is independent of the choice of path. Moreover we have

$$-\sqrt{-1} \partial \bar{\partial} \text{BC}_\Phi(H, H_0) = \Phi(H) - \Phi(H_0),$$

which can be checked by differentiating with respect to some parameter.

We will restrict to the case of Chern classes of holomorphic vector bundles. The Chern-Weil form for the metric H representing the k th Chern class of E is given by

$$c_k(H) = \frac{1}{k!} \left(\frac{\sqrt{-1}}{2\pi} \right)^k \sum_{\pi \in \Sigma_k} \text{sgn}(\pi) F_{\alpha_{\pi(1)}}^{\alpha_1} \wedge F_{\alpha_{\pi(2)}}^{\alpha_2} \wedge \dots \wedge F_{\alpha_{\pi(k)}}^{\alpha_k},$$

where $F = F_H$ and the usual summation convention is being used.

The Bott-Chern form for the k th Chern class is then given by

$$\begin{aligned} & \text{BC}_k(H, H_0) \\ &= -k\sqrt{-1} \int_0^1 \frac{1}{k!} \left(\frac{\sqrt{-1}}{2\pi} \right)^k \sum_{\pi \in \Sigma_k} \text{sgn}(\pi) (\dot{H}_t H_t^{-1})_{\alpha_{\pi(1)}}^{\alpha_1} F_{\alpha_{\pi(2)}}^{\alpha_2} \wedge \dots \wedge F_{\alpha_{\pi(k)}}^{\alpha_k} dt, \end{aligned}$$

where $F = F_{H_t}$. It satisfies

$$c_k(H) - c_k(H_0) = -\sqrt{-1}\partial\bar{\partial}\text{BC}_k(H, H_0).$$

Now consider the case where the vector bundle is $T'M$, the holomorphic tangent bundle of M . Then $g_{i\bar{j}}$, the Kähler metric on M , is a Hermitian metric on $T'M$. Let ϕ be in $P(M, \omega)$. Write g_ϕ for $g_{i\bar{j}} + \partial_i\partial_{\bar{j}}\phi$ and $c_k(\phi)$ for $c_k(g_\phi)$. Then, in normal coordinates for g_ϕ , we have

$$c_k(\phi) = \frac{1}{k!} \left(\frac{\sqrt{-1}}{2\pi} \right)^k \sum_{\pi \in \Sigma_k} \text{sgn}(\pi) F_{\alpha_{\pi(1)}\bar{\alpha}_1} \wedge \dots \wedge F_{\alpha_{\pi(k)}\bar{\alpha}_k},$$

writing $F_{\alpha\bar{\beta}}$ for $R_{\alpha\bar{\beta}i\bar{j}}dz^i \wedge d\bar{z}^{\bar{j}}$, where $R_{\alpha\bar{\beta}i\bar{j}}$ is the curvature tensor for g_ϕ .

3. Higher K-energy functionals

In this section we give an alternative proof of Theorem 1. Let $\{\phi_t\}_{0 \leq t \leq 1}$ be a smooth path in $P(M, \omega)$ with $\phi_0 = \phi_1$, and let λ be any constant. Calculate

$$\begin{aligned} & \int_0^1 \int_M \dot{\phi}_t (c_k(\phi_t) - \lambda \omega_t^k) \wedge \omega_t^{n-k} dt \\ &= \int_0^1 \int_M \dot{\phi}_t (c_k(\phi_t) - c_k(\phi_0) + c_k(\phi_0) - \lambda \omega_t^k) \wedge \omega_t^{n-k} dt \\ &= \int_0^1 \int_M \dot{\phi}_t (-\sqrt{-1}\partial\bar{\partial}\text{BC}_k(\phi_t, \phi_0) + c_k(\phi_0) - \lambda \omega_t^k) \wedge \omega_t^{n-k} dt \\ &= \int_0^1 \int_M (-\sqrt{-1}\partial\bar{\partial}\dot{\phi}_t) \wedge \text{BC}_k(\phi_t, \phi_0) \wedge \omega_t^{n-k} dt \\ &\quad + \int_0^1 \int_M \dot{\phi}_t (c_k(\phi_0) - \lambda \omega_t^k) \wedge \omega_t^{n-k} dt \\ &= -\frac{1}{n-k+1} \int_0^1 \int_M \text{BC}_k(\phi_t, \phi_0) \wedge \frac{\partial}{\partial t} (\omega_t^{n-k+1}) dt \\ &\quad + \int_0^1 \int_M \dot{\phi}_t (c_k(\phi_0) - \lambda \omega_t^k) \wedge \omega_t^{n-k} dt \\ &= \frac{1}{n-k+1} \int_0^1 \int_M \frac{\partial}{\partial t} \text{BC}_k(\phi_t, \phi_0) \wedge \omega_t^{n-k+1} dt \\ &\quad + \int_0^1 \int_M \dot{\phi}_t (c_k(\phi_0) - \lambda \omega_t^k) \wedge \omega_t^{n-k} dt. \end{aligned}$$

The next two lemmas will show that this expression is zero. The proof of the theorem will then follow immediately.

Lemma 3.1 *With $\{\phi_t\}$ as above,*

$$\int_0^1 \int_M \dot{\phi}_t(c_k(\phi_0) - \lambda \omega_t^k) \wedge \omega_t^{n-k} dt = 0.$$

Proof The argument is almost identical to one given in [13], but we will include it for the reader's convenience. Define

$$\phi_{t,s} = s\phi_t, \quad \text{and} \quad \omega_{t,s} = \omega + \sqrt{-1}\partial\bar{\partial}\phi_{t,s},$$

$$D(s) = \int_0^1 \int_M \frac{\partial \phi_{t,s}}{\partial t} (c_k(\phi_0) - \lambda \omega_{t,s}^k) \wedge \omega_{t,s}^{n-k} dt.$$

Notice that $D(0) = 0$. We will show that $D'(s) = 0$.

$$\begin{aligned} D'(s) &= \int_0^1 \int_M \frac{\partial^2 \phi_{t,s}}{\partial t \partial s} (c_k(\phi_0) - \lambda \omega_{t,s}^k) \wedge \omega_{t,s}^{n-k} dt \\ &\quad + \int_0^1 \int_M \frac{\partial \phi_{t,s}}{\partial t} ((n-k)c_k(\phi_0)\omega_{t,s}^{n-k-1} - \lambda n \omega_{t,s}^{n-1}) \wedge \sqrt{-1}\partial\bar{\partial}(\frac{\partial \phi_{t,s}}{\partial t}) dt \\ &= \int_0^1 \int_M \frac{\partial}{\partial t} \left(\frac{\partial \phi_{t,s}}{\partial s} \right) (c_k(\phi_0) - \lambda \omega_{t,s}^k) \wedge \omega_{t,s}^{n-k} dt \\ &\quad + \int_0^1 \int_M \frac{\partial \phi_{t,s}}{\partial s} ((n-k)c_k(\phi_0)\omega_{t,s}^{n-k-1} - \lambda n \omega_{t,s}^{n-1}) \wedge \sqrt{-1}\partial\bar{\partial}(\frac{\partial \phi_{t,s}}{\partial t}) dt \\ &= \int_0^1 \int_M \frac{\partial}{\partial t} \left(\frac{\partial \phi_{t,s}}{\partial s} (c_k(\phi_0) - \lambda \omega_{t,s}^k) \wedge \omega_{t,s}^{n-k} \right) dt \\ &= 0. \end{aligned}$$

Lemma 3.2 *With $\{\phi_t\}$ as above,*

$$\int_M \frac{\partial}{\partial t} (\text{BC}_k(\phi_t, \phi_0)) \wedge \omega_t^{n-k+1} = 0.$$

Proof Writing g for g_{ϕ_t} we have

$$\dot{g}_{i\bar{j}} g^{\bar{j}k} = (\partial_i \partial_{\bar{j}} \dot{\phi}_t) g^{\bar{j}k}.$$

Hence, using the formula for the Bott-Chern form, and working in normal coordinates,

$$\int_M \frac{\partial}{\partial t} \text{BC}_k(\phi_t, \phi_0) \wedge \omega_t^{n-k+1}$$

$$\begin{aligned}
&= \int_M \frac{-\sqrt{-1}}{(k-1)!} \left(\frac{\sqrt{-1}}{2\pi} \right)^k \sum_{\pi \in \Sigma_k} \text{sgn}(\pi) \partial_{\alpha_{\pi(1)}} \partial_{\bar{\alpha}_1} \dot{\phi}_t F_{\alpha_{\pi(2)} \bar{\alpha}_2} \wedge \dots \\
&\quad \dots \wedge F_{\alpha_{\pi(k)} \bar{\alpha}_k} \wedge \omega_t^{n-k+1} \\
&= \frac{-\sqrt{-1}}{(k-1)!} \left(\frac{\sqrt{-1}}{2\pi} \right)^k \int_M \sum_{\pi \in \Sigma_k} \text{sgn}(\pi) \nabla_{\alpha_{\pi(1)}} \nabla_{\bar{\alpha}_1} \dot{\phi}_t F_{\alpha_{\pi(2)} \bar{\alpha}_1} \wedge \dots \\
&\quad \dots \wedge F_{\alpha_{\pi(k)} \bar{\alpha}_k} \wedge \omega_t^{n-k+1}. \tag{1}
\end{aligned}$$

Integrating by parts, using the Bianchi identity, and defining τ to be the transposition in Σ_k which interchanges 1 and 2, we see that one term in the integrand of (1) will be

$$\begin{aligned}
& - \sum_{\pi \in \Sigma_k} \text{sgn}(\pi) \nabla_{\bar{\alpha}_1} \dot{\phi}_t \nabla_{\alpha_{\pi(1)}} F_{\alpha_{\pi(2)} \bar{\alpha}_2} \wedge F_{\alpha_{\pi(3)} \bar{\alpha}_3} \wedge \dots \wedge F_{\alpha_{\pi(k)} \bar{\alpha}_k} \wedge \omega_t^{n-k+1} \\
&= - \sum_{\pi \in \Sigma_k} \text{sgn}(\pi) \nabla_{\bar{\alpha}_1} \dot{\phi}_t \nabla_{\alpha_{\pi(2)}} F_{\alpha_{\pi(1)} \bar{\alpha}_2} \wedge F_{\alpha_{\pi(3)} \bar{\alpha}_3} \wedge \dots \wedge F_{\alpha_{\pi(k)} \bar{\alpha}_k} \wedge \omega_t^{n-k+1} \\
&= - \sum_{\pi \in \Sigma_k} \text{sgn}(\pi\tau) \nabla_{\bar{\alpha}_1} \dot{\phi}_t \nabla_{\alpha_{\pi(1)}} F_{\alpha_{\pi(2)} \bar{\alpha}_2} \wedge F_{\alpha_{\pi(3)} \bar{\alpha}_3} \wedge \dots \wedge F_{\alpha_{\pi(k)} \bar{\alpha}_k} \wedge \omega_t^{n-k+1} \\
&= \sum_{\pi \in \Sigma_k} \text{sgn}(\pi) \nabla_{\bar{\alpha}_1} \dot{\phi}_t \nabla_{\alpha_{\pi(1)}} F_{\alpha_{\pi(2)} \bar{\alpha}_2} \wedge F_{\alpha_{\pi(3)} \bar{\alpha}_3} \wedge \dots \wedge F_{\alpha_{\pi(k)} \bar{\alpha}_k} \wedge \omega_t^{n-k+1} \\
&= 0.
\end{aligned}$$

Every other term is zero by the same argument, and the lemma is proved.

The proof of Theorem 1 is now complete.

Remark 3.3 $M_{k,\omega}$ satisfies a cocycle condition. Namely, if ω' is another Kähler metric with $\omega' = \omega + \sqrt{-1}\partial\bar{\partial}\psi$, then

$$M_{k,\omega}(\phi) - M_{k,\omega'}(\phi - \psi) = M_{k,\omega}(\psi).$$

Remark 3.4 The critical points of $M_{k,\omega}$ are metrics ω with $\Lambda^k c_k(\omega)$ constant. An example of such a metric is the Fubini-Study metric on $\mathbf{CP}^n$.

4. Higher Futaki invariants

In this section we give a proof of Theorem 2. Let X be a holomorphic vector field. Recall that the Lie derivative of X on forms is given by

$$\mathcal{L}_X = i_X \circ d + d \circ i_X.$$

The interior product $i_X\omega$ is a $(0,1)$ form which is $\bar{\partial}$ -closed since ω is Kähler. Hence there exists a smooth function θ_X and a harmonic $(0,1)$ form α such that

$$i_X\omega = \alpha - \bar{\partial}\theta_X.$$

We have the following lemma.

Lemma 4.1

$$\mathcal{F}_{k,\omega}(X) = -(n-k+1)\sqrt{-1} \int_M \theta_X(c_k(\omega) - Hc_k(\omega)) \wedge \omega^{n-k},$$

where θ_X is as given above.

Proof Integrating by parts,

$$\begin{aligned} \mathcal{F}_{k,\omega}(X) &= \int_M \mathcal{L}_X f_k \wedge \omega^{n-k+1} \\ &= - \int_M f_k \wedge \mathcal{L}_X \omega^{n-k+1} \\ &= - \int_M f_k \wedge \partial i_X \omega^{n-k+1} \\ &= -(n-k+1) \int_M f_k \wedge \partial(\alpha - \bar{\partial}\theta_X) \wedge \omega^{n-k} \\ &= (n-k+1) \int_M f_k \wedge \partial\bar{\partial}\theta_X \wedge \omega^{n-k} \\ &= -(n-k+1)\sqrt{-1} \int_M \theta_X(c_k(\omega) - Hc_k(\omega)) \wedge \omega^{n-k}. \end{aligned}$$

To finish the proof of the theorem, let Φ_t be the integral curve of X_R as before, and let ϕ_t be given by

$$\Phi_t^*\omega = \omega_t = \omega + \sqrt{-1}\partial\bar{\partial}\phi_t, \quad \int_M \phi_t \omega^n = 0.$$

Then

$$\mathcal{L}_{X_R}\omega_t = \sqrt{-1}\partial\bar{\partial}\dot{\phi}_t.$$

Hence

$$\partial i_X \omega_t + \bar{\partial} i_{\bar{X}} \omega_t = \frac{1}{2}\sqrt{-1}(\partial\bar{\partial}\dot{\phi}_t - \bar{\partial}\partial\dot{\phi}_t)$$

and so

$$i_X \omega_t - \frac{1}{2}\sqrt{-1}\partial\bar{\partial}\dot{\phi}_t = \alpha_t,$$

where α_t is a $(0,1)$ form satisfying $\text{Re}(\partial\alpha_t) = 0$. It follows from above that

$$\begin{aligned} 2\text{Re}(\mathcal{F}_{k,\omega_t}(X)) &= -(n-k+1) \int_M \dot{\phi}_t(c_k(\omega_t) - Hc_k(\omega_t)) \wedge \omega_t^{n-k} \\ &= -(n-k+1) \int_M \dot{\phi}_t(c_k(\omega_t) - \mu_k \omega_t^k) \wedge \omega_t^{n-k} \\ &= \frac{d}{dt} M_{k,\omega}(\phi_t). \end{aligned}$$

Finally we must show that $\mathcal{F}_{k,\omega_t}(X) = \mathcal{F}_{k,\omega}(X)$. This is immediate from [1], or alternatively can be seen as follows. Since

$$\mathcal{L}_{X_R} \omega_t = \Phi_t^* \mathcal{L}_{X_R} \omega,$$

we have

$$\partial\bar{\partial}\dot{\phi}_t = \partial\bar{\partial}\Phi_t^* \dot{\phi}_0,$$

and so $\dot{\phi}_t = \Phi_t^* \dot{\phi}_0$. Hence

$$\begin{aligned} \int_M \dot{\phi}_t(c_k(\omega_t) - \mu_k \omega_t^k) \wedge \omega_t^{n-k} &= \int_M \Phi_t^* \dot{\phi}_0(\Phi_t^* c_k(\omega) - \mu_k \Phi_t^* \omega^k) \wedge \Phi_t^* \omega^{n-k} \\ &= \int_M \dot{\phi}_0(c_k(\omega) - \mu_k \omega^k) \wedge \omega^{n-k}. \end{aligned}$$

5. Higher K-energy functionals and Donaldson's Lagrangian

In this section we give a proof of Theorem 3 and give two alternative formulas for the higher K-energy functionals. It is pointed out in [15] (see also [5] and [13]) that the K-energy can be written without a path integral as

$$\begin{aligned} M_{1,\omega}(\phi) &= \int_M \frac{1}{2\pi} \log \left(\frac{\omega_\phi^n}{\omega^n} \right) \omega_\phi^n \\ &\quad - \int_M \phi \left(\frac{\sqrt{-1}}{2\pi} \text{Ric}(\omega) \wedge \sum_{i=0}^{n-1} \omega^i \wedge \omega_\phi^{n-1-i} - \frac{\mu_1}{n+1} \sum_{i=0}^n \int_M \phi \omega^i \wedge \omega_\phi^{n-i} \right). \end{aligned}$$

This appears to be both a natural and useful expression for the K-energy (see [12]). Notice that

$$c_1(\omega) = \frac{\sqrt{-1}}{2\pi} \text{Ric}(\omega) \quad \text{and} \quad \frac{1}{2\pi} \log \left(\frac{\omega_\phi^n}{\omega^n} \right) = \text{BC}_1(\phi, 0),$$

where to be precise, the above is only a *representative* of $\text{BC}_1(\phi, 0)$. The following lemma generalizes the above formula to the higher K-energy functionals.

Lemma 5.1

$$M_{k,\omega}(\phi) = \int_M BC_k(\phi, 0) \wedge \omega_\phi^{n-k+1} \\ - \int_M \phi \left(c_k(0) \wedge \sum_{i=0}^{n-k} \omega^i \wedge \omega_\phi^{n-k-i} - \frac{(n-k+1)}{n+1} \mu_k \sum_{i=0}^n \omega^i \wedge \omega_\phi^{n-i} \right).$$

Proof For a path $\{\phi_t\}$ in $P(M, \omega)$, calculate

$$\begin{aligned} \int_M \frac{\partial}{\partial t} (BC_k(\phi_t, 0) \wedge \omega_t^{n-k+1}) \\ = \int_M \frac{\partial}{\partial t} (BC_k(\phi_t, 0)) \wedge \omega_t^{n-k+1} \\ + (n-k+1) \int_M BC_k(\phi_t, 0) \wedge \omega_t^{n-k} \wedge \sqrt{-1} \partial \bar{\partial} \dot{\phi}_t \\ = -(n-k+1) \int_M \dot{\phi}_t (c_k(\phi_t) - c_k(0)) \wedge \omega_t^{n-k}, \end{aligned} \quad (2)$$

using Lemma 3.2 for the first term and integrating by parts for the second. Also

$$\begin{aligned} \int_M \frac{\partial}{\partial t} (\phi_t c_k(0) \wedge \sum_{i=0}^{n-k} \omega^i \wedge \omega_t^{n-k-i}) \\ = \int_M \dot{\phi}_t c_k(0) \wedge \sum_{i=0}^{n-k} \omega^i \wedge \omega_t^{n-k-i} \\ + \int_M \phi_t c_k(0) \wedge \sum_{i=0}^{n-k} (n-k-i) \omega^i \wedge \omega_t^{n-k-i-1} \wedge \sqrt{-1} \partial \bar{\partial} \dot{\phi}_t \\ = \int_M \dot{\phi}_t c_k(0) \wedge \sum_{i=0}^{n-k} \omega^i \wedge \omega_t^{n-k-i} \\ + \int_M \dot{\phi}_t c_k(0) \wedge \sum_{i=0}^{n-k} (n-k-i) \omega^i \wedge \omega_t^{n-k-i-1} \wedge (\omega_t - \omega) \\ = \int_M \dot{\phi}_t c_k(0) \wedge \left(\sum_{i=0}^{n-k} (n-k-i+1) \omega^i \wedge \omega_t^{n-k-i} \right. \\ \left. - (n-k-i) \omega^{i+1} \wedge \omega_t^{n-k-i-1} \right) \\ = (n-k+1) \int_M \dot{\phi}_t c_k(0) \wedge \omega_t^{n-k}. \end{aligned} \quad (3)$$

Similarly,

$$\int_M \frac{\partial}{\partial t} \left(\frac{(n-k+1)\mu_k}{n+1} \phi_t \sum_{i=0}^n \omega^i \wedge \omega_t^{n-i} \right) = (n-k+1)\mu_k \int_M \dot{\phi}_t \omega_t^n. \quad (4)$$

The lemma follows from (2), (3) and (4).

We will need a slightly different formula for $M_{k,\omega}$.

Corollary 5.2

$$\begin{aligned} M_{k,\omega}(\phi) &= \int_M BC_k(\phi, 0) \wedge \omega^{n-k+1} \\ &\quad - \int_M \phi \left(c_k(\phi) \wedge \sum_{i=0}^{n-k} \omega^i \wedge \omega_\phi^{n-k-i} - \frac{(n-k+1)}{n+1} \mu_k \sum_{i=0}^n \omega^i \wedge \omega_\phi^{n-i} \right). \end{aligned}$$

Proof

$$\begin{aligned} &\int_M BC_k(\phi, 0) \wedge \omega_\phi^{n-k+1} \\ &= \int_M BC_k(\phi, 0) \wedge (\omega + \sqrt{-1} \partial \bar{\partial} \phi)^{n-k+1} \\ &= \int_M BC_k(\phi, 0) \wedge \sum_{i=0}^{n-k+1} \binom{n-k+1}{i} \omega^i \wedge (\omega_\phi - \omega)^{n-k+1-i} \\ &= \int_M BC_k(\phi, 0) \wedge \omega^{n-k+1} \\ &\quad + \int_M BC_k(\phi, 0) \wedge \sqrt{-1} \partial \bar{\partial} \phi \wedge \sum_{i=0}^{n-k} \binom{n-k+1}{i} \omega^i \wedge (\omega_\phi - \omega)^{n-k-i} \\ &= \int_M BC_k(\phi, 0) \wedge \omega^{n-k+1} - \int_M \phi (c_k(\phi) - c_k(0)) \wedge \sum_{i=0}^{n-k} \omega^i \wedge \omega_\phi^{n-k-i}, \end{aligned}$$

using, for the last line, the elementary identity

$$\sum_{i=0}^m \binom{m+1}{i} x^i (y-x)^{m-i} = \sum_{i=0}^m x^i y^{m-i},$$

which can be seen by observing that each side is equal to

$$(y^{m+1} - x^{m+1})/(y-x).$$

The corollary now follows immediately from Lemma 5.1.

We now prove Theorem 3. First note that we have

$$\begin{aligned} \text{BC}_1(\phi, 0) &= \frac{1}{2\pi} \int_0^1 \text{Tr}(\dot{g}_t g_t^{-1}) dt, \quad \text{and} \\ \text{BC}_2(\phi, 0) &= \frac{\sqrt{-1}}{(2\pi)^2} \int_0^1 \left(\text{Tr}(\dot{g}_t g_t^{-1}) \text{Tr}(F_t) - \text{Tr}(\dot{g}_t g_t^{-1} F_t) \right) dt, \end{aligned}$$

where F_t is the curvature of g_t . Then using the fact that $\lambda = 2\pi\mu_1$ we have

$$\begin{aligned} L(\phi) &= n(2\pi)^2 \int_M \int_0^1 \frac{\sqrt{-1}}{(2\pi)^2} \text{Tr}(\dot{g}_t g_t^{-1} F_t) \wedge \omega^{n-1} dt \\ &\quad - (2\pi)^2 \mu_1 \int_M \int_0^1 \frac{1}{2\pi} \text{Tr}(\dot{g}_t g_t^{-1}) \omega^n dt \\ &= -n(2\pi)^2 \int_M \text{BC}_2(\phi, 0) \wedge \omega^{n-1} - (2\pi)^2 \mu_1 \int_M \text{BC}_1(\phi, 0) \omega^n \\ &\quad + n\sqrt{-1} \int_M \int_0^1 \text{Tr}(\dot{g}_t g_t^{-1}) \text{Tr}(F_t) \wedge \omega^{n-1} dt. \end{aligned} \tag{5}$$

The last term is equal to

$$\begin{aligned} &n\sqrt{-1} \int_M \int_0^1 (\Delta_t \dot{\phi}_t) \text{Ric}(\omega_t) \wedge \omega^{n-1} dt \\ &= \frac{n\sqrt{-1}}{2} \int_M \log \left(\frac{\omega_\phi^n}{\omega^n} \right) (\text{Ric}(\omega_t) + \text{Ric}(\omega)) \wedge \omega^{n-1} \\ &= n\pi \int_M \log \left(\frac{\omega_\phi^n}{\omega^n} \right) (c_1(\phi) + c_1(0)) \wedge \omega^{n-1}, \end{aligned} \tag{6}$$

where a straightforward integration by parts has been used for the second line. The proof of the theorem now follows from Corollary 5.2 and equations (5) and (6).

References

- [1] Bando, S. *An obstruction for Chern class forms to be harmonic*, unpublished.
- [2] Bando, S., Mabuchi, T. *Uniqueness of Einstein Kähler metrics modulo connected group actions* in: Oda, T. (ed.) *Algebraic Geometry*, Sendai 1985. (Adv. Stud. Pure Math., Vol 10) Amsterdam: North-Holland & Tokyo: Kinokuniya (1987)

- [3] Bando, S., Mabuchi, T. *On some integral invariants on complex manifolds. I*, Proc. Japan Acad., 62 (1986), 197-200
- [4] Bott, R. and Chern, S.S. *Hermitian vector bundles and the equidistribution of the zeroes of their holomorphic sections*, Acta Math., 114 (1965), 71-112
- [5] Chen, X. X. *On the lower bound of the Mabuchi energy and its application*, Int. Math Res. Notices, 12 (2000), 607-623
- [6] Donaldson, S.K. *Anti self-dual Yang-Mills connections over complex algebraic surfaces and stable vector bundles*, Proc. London Math. Soc. (3), 50 (1985), 1-26
- [7] Donaldson, S.K. *Infinite determinants, stable bundles and curvature*, Duke Math. Journ., 54, No. 1 (1987), 231-247
- [8] Futaki, A. *An obstruction to the existence of Einstein Kähler metrics*, Invent. Math. 73 (1983), 437-443
- [9] Futaki, A., *Kähler-Einstein metrics and integral invariants*, Lecture Notes in Mathematics 1314, Springer-Verlag, Berlin Heidelberg (1988)
- [10] Futaki, A., Mabuchi, T. and Sakane, Y. *Einstein-Kähler metrics with positive Ricci curvature*, in Kähler Metrics and Moduli Spaces (Adv. Studies in Pure Math., Vol 18, II) (1990), 11-83
- [11] Mabuchi, T. *K-energy maps integrating Futaki invariants*, Tôhoku Math. Journ., 38 (1986), 575-593
- [12] Phong, D.H. and Sturm, J. *Stability, energy functionals and Kähler-Einstein metrics*, preprint, mathDG.0203254
- [13] Tian, G. *The K-energy on hypersurfaces and stability*, Comm. Anal. and Geom., Vol 2., No. 2 (1994), 239-265
- [14] Tian, G. *Kähler-Einstein metrics with positive Ricci curvature*, Inv. Math. 130 (1997), 1-39
- [15] Tian, G. *Canonical metrics in Kähler geometry*, Lectures in Mathematics ETH Zürich, Birkhäuser Verlag, Basel, Boston, Berlin, 2000

- [16] Tian, G. *Bott-Chern forms and geometric stability*, Discrete Contin. Dynam. Systems, 6 (2000), 211-220
- [17] Yau, S. T. *Open problems in geometry*, Proc. Symposia Pure Math. 54 (1993), 1-28
- [18] Yau, S. T. *Review of Kähler-Einstein metrics in Algebraic Geometry*, Israel Math. Conf. Proceedings, 9 (1996), 433-443